



A service of the National Library of Medicine  
and the National Institutes of Health

My NCBI  
[Sign In] [Register]

All Databases PubMed Nucleotide Protein Genome Structure OMIM PMC Journals Books

Search PubMed for Go Clear

Limits Preview/Index History Clipboard Details

About Entrez

Display Citation Show 20 Sort by Send to

Text Version

All: 1 Review: 0

## Entrez PubMed

Overview  
Help | FAQ  
Tutorials

New/Noteworthy  
E-Utilities

## PubMed Services

Journals Database  
MeSH Database  
Single Citation Matcher  
Batch Citation Matcher  
Clinical Queries  
Special Queries  
LinkOut  
My NCBI

## Related Resources

Order Documents  
NLM Mobile  
NLM Catalog  
NLM Gateway  
TOXNET  
Consumer Health  
Clinical Alerts  
ClinicalTrials.gov  
PubMed Central

1: Int J Dermatol. 1998 Mar;37(3):211-4.

Related Articles, Links



## Familial pemphigus vulgaris in mother and daughter.

Starzycki Z, Chorzelski TP, Jablonska S.

Department of Dermatology, Collegium Medicum Jagiellonian University,  
Cracow, Poland.

**BACKGROUND:** Pemphigus vulgaris is an autoimmune disease in which genetics appears to be of basic importance. Although association with certain human leukocyte antigen (HLA) alleles has been found in some ethnic groups and individuals, no true disease susceptibility genes have been established, and familial cases are very unusual. **METHODS:** We report a Polish family with pemphigus vulgaris in the mother and daughter. The diagnosis was confirmed by cytologic, histologic, and immunofluorescence studies. **RESULTS:** The course was severe and the disease long-lasting in the mother, probably due to treatment with small doses of corticosteroids without immunosuppressive drugs. In the daughter, treated with larger doses of corticosteroids and azathioprine, the lesions regressed within 4 months, after which maintenance therapy was instituted with 10 mg of prednisone daily. The HLA studies performed in the daughter and her three children after the mother had died showed identical haplotypes in both the patient and the healthy children. The patient has given birth to a healthy child while still having a high titer of intercellular (IC) antibodies. **CONCLUSIONS:** The familial occurrence of pemphigus in first-degree relatives is suggestive of inherited susceptibility to the disease, transmitted as a dominant trait. The identical haplotypes in the healthy children of the patient favor the role of other, unknown factors required for the development of the disease in predisposed individuals.

## Publication Types:

- Case Reports

## MeSH Terms:

- Adult
- Aged
- Diagnosis, Differential

- [Female](#)
- [HLA Antigens/genetics](#)
- [Haplotypes](#)
- [Humans](#)
- [Pemphigus, Benign Familial/diagnosis\\*](#)
- [Pemphigus, Benign Familial/drug therapy](#)
- [Pemphigus, Benign Familial/genetics\\*](#)
- [Pemphigus, Benign Familial/immunology](#)

Substances:

- [HLA Antigens](#)

PMID: 9556111 [PubMed - indexed for MEDLINE]

Display	<input type="text" value="Citation"/>	Show	<input type="text" value="20"/>	Sort by	<input type="text" value=""/>	Send to	<input type="text" value=""/>
---------	---------------------------------------	------	---------------------------------	---------	-------------------------------	---------	-------------------------------

[Write to the Help Desk](#)  
[NCBI](#) | [NLM](#) | [NIH](#)  
[Department of Health & Human Services](#)  
[Privacy Statement](#) | [Freedom of Information Act](#) | [Disclaimer](#)